



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Putting $x = \cos\theta + i\sin\theta$, and employing the formula $(\cos\theta + i\sin\theta)^m = \cos m\theta + i\sin m\theta$, we obtain after putting the real parts of both members equal, and making all necessary reductions, for the sum of the second series

$$= \frac{\cos\theta - n\cos n\theta + (n-1)\cos(n+1)\theta}{4\sin^2 \frac{1}{2}\theta};$$

so that the sum of the given series

$$= \frac{n\cos \frac{1}{2}(n+1)\theta \sin \frac{1}{2}n\theta}{\sin \frac{1}{2}\theta} + \frac{\cos\theta - n\cos n\theta + (n-1)\cos(n+1)\theta}{4\sin^2 \frac{1}{2}\theta}.$$

To test this formula we must of course, leave the coefficient n of the first expression unchanged, while in all the other factors and terms which involve n , n must be put successively $= 1, 2, 3, 4$, etc.

Also solved by *E. W. MORRELL*.

69. Proposed by *C. E. WHITE*, A. M., Trafalgar, Indiana.

Prove that $x^n \pm x^{n-1} + x^{n-2} \pm \dots + (\pm 1)^{n-1}x + (\pm 1)^n = (x \mp 1)^n \pm A(x \mp 1)^{n-1} + B(x \mp 1)^{n-2} \pm \dots + (\pm 1)^n x$, where A, B, C, \dots are the binomial coefficients of the $(n+1)$ th order.

I. Solution by *O. W. ANTHONY*, M. Sc., Professor of Mathematics in Columbian University, Washington, D. C.

$$x^n \pm x^{n-1} + x^{n-2} + \dots, \text{ etc.} = (x^{n+1} - 1)/(x - 1) \dots \dots \dots (1),$$

$$\text{or } x^{n+1} + 1)/(x + 1) \dots \dots \dots (2), = \{[(x - 1) + 1]^{n+1} - 1\}/(x - 1), \text{ or}$$

$$\{[(x + 1) - 1]^{n+1} + 1\}/(x + 1), = (x - 1)^n + C_{n+1}^2(x - 1)^{n-1} + C_{n+1}^3(x - 1)^{n-2} + \dots,$$

$$\text{or } (x + 1)^n - C_{n+1}^2(x + 1)^{n-1} + C_{n+1}^3(x - 1)^{n-2} - \dots,$$

$$= (x \mp 1)^n \pm A(x \mp 1)^{n-1} + B(x \mp 1)^{n-2} + \dots$$

II. Solution by *E. W. MORRELL*, Professor of Mathematics in Montpelier Seminary, Montpelier, Vermont.

$$\text{Let } K = x^n \pm x^{n-1} + x^{n-2} \pm \dots + (\pm 1)^{n-1}x + (\pm 1)^n.$$

Put $x = y \pm 1$, expanding and observing that the sign of the last term of each expression is \pm if n is odd but $+$ if n is even, we may write :

$$x^n = (y \pm 1)^n = y^n \pm n y^{n-1} + \frac{1}{2}[n(n-2)]y^{n-2} \pm \dots + (\pm 1)^{n-1}ny + (\pm 1)^n$$

$$\pm x^{n-1} = \pm (y \pm 1)^{n-1} = \pm y^{n-1} + (n-1)y^{n-2} \pm \dots + (\pm 1)^{n-1}(n-1)y + (\pm 1)^n$$

$$x^{n-2} = (y \pm 1)^{n-2} = \dots y^{n-2} \pm \dots + (\pm 1)^{n-1}(n-2)y + (\pm 1)^n$$

$$\text{etc.} \dots \dots \dots \text{etc.}$$

$$(\pm 1)^{n-1}x = (\pm 1)^{n-1}(y \pm 1) = \dots (\pm 1)^{n-1}y + (\pm 1)^n$$

$$(\pm 1)^n = \dots (\pm 1)^n.$$

By adding, and simplifying the coefficients of y , we have

$$K = y^n \pm (n+1)y^{n-1} + \frac{1}{2}[(n+1)n]y^{n-2} \pm \dots + (\pm 1)^{n-1} \frac{1}{2}[(n+1)n]y + (\pm 1)^n(n+1),$$

which has binomial coefficients of the $(n+1)$ th order. Substituting A , B , C ,
..... for the coefficients and restoring the values of y ,

$$K = (x \mp 1)^n \pm A(x \mp 1)^{n-1} + B(x \mp 1)^{n-2} \pm \dots + (\pm 1)^{n-1}B(x \mp 1) + (\pm 1)^n A.$$

[Expanding and combining the terms of the second member, we get the first member for a result. ZERR.]

GEOMETRY.

Conducted by B. F. FINKEL, Springfield, Mo. All contributions to this department should be sent to him.

SOLUTIONS OF PROBLEMS.

43. Proposed by J. F. W. SCHEFFER, A. M., Hagerstown, Maryland.

The consecutive sides of a quadrilateral are a , b , c , d . Supposing its diagonals to be equal, find them and also the area of the quadrilateral.

II. Solution by A. H. BELL, Hillsboro, Illinois.

The solution as published simply demonstrates this theorem, $a^2 + b^2 + c^2 + d^2 = 2x^2 + 4\overline{IK}^2$, with two unknowns, and is then solved for a particular case.

Let the sides of the quadrilateral AB , BC , CD , and AD , be a , b , c , and d ; and the diagonals each $= 2x$; $x+y$, $x-y$ = the segments AO and OC ; and BO and OD = $x+z$ and $x-z$. In the triangles AOB , BOC , and COD we have $(x+y)\cos A + (x+z)\cos B = a$, $\cos A = (4x^2 + a^2 - b^2)/(4ax)$, $\cos B = (4x^2 + a^2 - d^2)/(4ax)$, making

